

The virtual scaling function of twist operators in the $\mathcal{N} = 6$ Chern-Simons theory

Matteo Beccaria and Guido Macorini

Physics Department, Salento University, Via Arnesano, 73100 Lecce
INFN, Sezione di Lecce
E-mail: matteo.beccaria@le.infn.it

ABSTRACT: We consider twist- L operators in the planar $\mathcal{N} = 6$ superconformal Chern-Simons ABJM theory. Their anomalous dimension $\gamma_L^{\text{CS}}(N)$ is a function of the twist L , the spin N , and the dressed coupling of ABJM. We show that at next-to-leading order in the large spin expansion, this anomalous dimension is related to that of $\mathcal{N} = 4$ SYM twist operators by a simple scaling law.

Contents

1. Introduction	1
2. Twist operators in ABJM and their exact anomalous dimensions	4
3. The fourth order twist-2 anomalous dimension	5
4. The virtual scaling function in ABJM from weak coupling	6
5. The NLO BES equation in $\mathcal{N} = 4$	7
6. Analysis of the ABJM $\mathfrak{sl}(2)$ Baxter equation	8
7. The integral equation determining $B_L(g)$	10
8. Comments	10
A. Next-to-leading large spin expansion of harmonic sums	11

1. Introduction

The maximally supersymmetric $\mathcal{N} = 4$ SYM theory is an ideal theoretical laboratory for the investigation of planar gauge theory. At weak coupling, all-order integrability leads to long range Bethe Ansatz equations which are an efficient computational tool and replace the usual diagrammatical expansion [1]. At strong coupling, AdS/CFT correspondence provides a non trivial reformulation in terms of type IIB superstring propagating on $AdS_5 \times S^5$ [2]. This leads to quantitative predictions which are hardly achievable by different means.

In this perspective, the so-called twist operators are an important probe of various universal quantities characterizing the gauge dynamics. At weak coupling, they are gauge invariant single trace composite operators belonging to a perturbatively closed $\mathfrak{sl}(2)$ sector. They are built as a matrix product of L fundamental $\mathcal{N} = 4$ SYM scalars φ with a certain number N of covariant derivatives spread along the chain, $\mathcal{O}_{L,N} \sim \text{Tr}(\varphi^{L-1} \mathcal{D}^N \varphi)$. The parameters L and N are called the twist and spin of the operator for clear reasons. The dual state at strong coupling is well identified in twist-2 and is described classically by a rotating folded string configuration [3, 4].

The analysis of twist operators is based on the study of their large spin limit, $N \rightarrow \infty$. In this regime, we can study their anomalous dimension $\gamma_{L,N}(g)$ as a function of the twist

L and, of course, the planar coupling g . The next-to-leading (NLO) expansion of $\gamma_{L,N}(g)$ has the general simple form [5, 6, 7, 8].

$$\gamma_{L,N}(g) = f(g)(\log N + \gamma_E - (L-2) \log 2) + B_L(g) + \dots, \quad (1.1)$$

where we have neglected terms vanishing at large N ¹. The coupling dependent coefficient of the logarithm is equal to twice the conventional QCD cusp anomaly [7] and is commonly referred as the *scaling function*. It is an universal quantity which is known at four loops by diagrammatical methods [14, 15]. An integral equation for $f(g)$ has been proposed in [16] (BES). It can generate the weak coupling expansion of $f(g)$ at any desired order. The strong coupling expansion of $f(g)$ is also known [17] (see also [18]).

From a physical point of view, the cusp anomaly defines what is called the physical coupling which measures the (universal) intensity of soft gluon radiation [10]. On the other hand, the physical meaning of the subleading constant term $B_L(g)$ is more subtle. It has been partially clarified in the recent analysis of Dixon-Magnea-Sterman [19] (DMS). From the work of DMS, we know that B_L is related to a piece of the infrared divergence of on-shell scattering amplitudes in $\mathcal{N} = 4$ SYM. The singular part of the amplitudes can be written schematically in the following factorized form [20]

$$\mathcal{A}_{\text{IR}} \sim \exp \left(\frac{f^{(-2)}(g)}{\varepsilon^2} - \frac{G^{(-1)}(g)}{\varepsilon} \right), \quad (1.2)$$

where ε is the IR dimensional regulator and $F^{(-n)}$ is the n -th times iterated logarithmic integral. The new function $G(g)$ is the collinear anomalous dimension [21, 22]. The important remark of DMS is that (at least for twist-2) G is the sum of a universal contribution G_{eik} and the subleading constant $B_2(g)$. The universal part is an eikonal contribution which can be extracted from the soft logarithms of the Drell-Yan process or from a suitable rectangular light-like Wilson loop. The non-universal piece $B_2(g)$ comes from the virtual contribution to the Altarelli-Parisi splitting kernel [23]. In Mellin space, this is precisely the subleading constant term appearing in Eq. (1.1). We shall call it the *virtual scaling function*. The DMS proposal has been recently confirmed at strong coupling in [24].

For the case of $\mathcal{N} = 4$ SYM, an integral equation for $B_L(g)$ has been analyzed in [25, 46]. The equation is derived neglecting wrapping effects. These are well known at leading order for $\mathfrak{sl}(2)$ twist operators [26]. At weak coupling, they do not affect the constant $B_L(g)$. Also at strong coupling, this turns out to be true as an outcome of the analysis of [25].

Recently, twist operators have also been introduced and studied in the ABJM theory [27] which has an integrable structure quite close to that of $\mathcal{N} = 4$ SYM, despite being a very different theory. ABJM is a 3d $U(N) \times U(N)$ Chern-Simons gauge theory with opposite levels $+k, -k$. It has $\mathcal{N} = 6$ superconformal symmetry and describes the low energy limit of N parallel M2-branes at a $\mathbb{C}^4/\mathbb{Z}_k$ singular point. For large N , k and fixed ratio

¹The subleading corrections to Eq. (1.1) are very interesting and are related to a generalized Gribov-Lipatov reciprocity as first discussed in [9, 10, 11]. It is a property very well tested at weak coupling [12], and partially investigated in string theory [13].

$\lambda = N/k$, we identify $Z_k \simeq S^1$ and M theory on the gravity dual orbifold $AdS_4 \times S^7/\mathbb{Z}_k$ reduces to type IIA string on $AdS_4 \times \mathbb{CP}^3$. The classical string integrability [28, 29, 30] has a gauge theory quantum counterpart first analyzed in [31, 32]. In [33], a set of all-loop Bethe-Ansatz equations have been proposed for the full $\mathfrak{osp}(2, 2|6)$ theory consistent with the string algebraic curve at strong coupling [34]. The equations depend on a dressed coupling $h(\lambda)$ which takes into account the fact that the one-magnon dispersion relation is not protected by supersymmetry [35, 36, 37, 38]. The conjectured S -matrix is worked out in [39].

ABJM twist operators and their anomalous dimensions are discussed in [33, 40, 41]. The similitudes with $\mathcal{N} = 4$ SYM are many. At strong coupling, the dual string state is a folded string rotating in AdS_3 with large spin N and with angular momentum $J \sim \log N$ in $\mathbb{CP}^3$ [42] not so different than the $AdS_5 \times S^5$ case. At weak coupling, their anomalous dimension is captured by all-loop Bethe Ansatz equation which are precisely those of $\mathcal{N} = 4$ SYM apart from a phase twist. This phase deformation [43] changes drastically some known fine features which are shared by $\mathcal{N} = 4$ SYM and QCD (Low-Burnett-Kroll wisdom, Gribov-Lipatov reciprocity). Nevertheless, the scaling function of ABJM is simply halved, apart from the unavoidable replacement $g \rightarrow h(g)$, and we have [33] (see also [44])

$$f^{\text{CS}}(h) = \frac{1}{2} f^{\mathcal{N}=4}(h), \quad (1.3)$$

where, here and in the following, CS stands for Chern-Simons and identifies the ABJM case. This factor $1/2$ can be understood in terms of the mode number change due to the phase twist in the (thermodynamical) large spin limit.

One immediately asks the following questions. What about the subleading constants appearing in Eq. (1.1)? How do they change in ABJM? Is there a simple relation with the $\mathcal{N} = 4$ SYM values? The result of this paper is that at next-to-leading order, *i.e.* at the level of the expansion Eq. (1.1), we can write

$$\gamma_L^{\text{CS}}(N) = \frac{1}{2} \gamma_{2L}^{\mathcal{N}=4}(2N), \quad (\text{NLO}). \quad (1.4)$$

In other words, we predict

$$\gamma_{L,N}^{\text{CS}}(h) = \frac{1}{2} f^{\mathcal{N}=4}(h) (\log(2N) + \gamma_E - 2(L-1) \log 2) + \frac{1}{2} B_{2L}^{\mathcal{N}=4}(h) + \dots \quad (1.5)$$

This conclusion is derived under the same hypothesis about wrapping we assumed in $\mathcal{N} = 4$ SYM². Notice that Eq. (1.4) is false beyond NLO order.


The plan of the paper is as follows. In Section 2, we briefly recall a few necessary basic definitions. In Section 3, we present a new closed formula for the (asymptotic) anomalous dimension of ABJM twist-2 operators at fourth order, *i.e.* eight loops. In Section 4, we compute and collect our results for the NLO large spin expansion of the ABJM anomalous dimensions. In Section 5, we briefly recall the form of the NLO BES equation in $\mathcal{N} = 4$

²A first positive test of this assumption has been presented in [41] for twist-1 and twist-2 operators at weak coupling and leading wrapping order.

SYM as a necessary preliminary steps for the introduction of its modified version valid in ABJM. Section 6 illustrates the analysis of the one-loop ABJM Baxter equation which describe twist operators. The analysis is aimed at providing arguments for the scaling behaviour of the hole solutions to the Bethe Ansatz equations. In Section 7, we report the $\mathcal{N} = 4$ and ABJM NLO BES equations in unified form and prove the scaling relation as a simple consequence. Finally, Section 8 is devoted to some final comments.

2. Twist operators in ABJM and their exact anomalous dimensions

The all-loop Bethe equations for ABJM has been proposed in [33]. They are associated with the $\mathfrak{osp}(2, 2|6)$ Dynkin diagram (in the fermionic $\eta = -1$ grading)


(2.1)

Twist operators in the $\mathfrak{sl}(2)$ sector are obtained by exciting the same number N of u_4 and $u_{\bar{4}}$ roots. As in the $\mathcal{N} = 4$ case, we shall refer to the integer L as the twist of the operator. More details can be found in [40].

Bethe Ansatz equations are written in terms of the deformed spectral parameters $x^\pm(u)$ defined by

$$x^\pm + \frac{1}{x^\pm} = \frac{1}{h} \left(u \pm \frac{i}{2} \right), \quad (2.2)$$

where $h(\lambda)$ is the interpolating coupling entering the one-magnon dispersion relation. For twist L operators they are

$$\left(\frac{x_k^+}{x_k^-} \right)^L = - \prod_{j \neq k}^N \frac{u_k - u_j + i}{u_k - u_j - i} \left(\frac{x_k^- - x_j^+}{x_k^+ - x_j^-} \right)^2 \sigma_{\text{BES}}^2. \quad (2.3)$$

The only difference compared with $\mathcal{N} = 4$ SYM is the extra minus sign whose effects are discussed in [38, 41]. The factor σ_{BES} is the Beisert-Eden-Staudacher dressing phase. The momentum constraint is automatically satisfied for Bethe root distributions symmetric under $u \rightarrow -u$ as those we are interested in.

The contribution to the energy/anomalous dimension from the Asymptotic Bethe Ansatz (ABA) equations is conveniently written in terms of

$$p(u) = -i \log \frac{x^+(u)}{x^-(u)}, \quad u(p) = \frac{1}{2} \cot \frac{p}{2} \sqrt{1 + 16 h^2 \sin^2 \frac{p}{2}}, \quad (2.4)$$

and reads

$$\gamma_L^{\text{CS}}(N, h) = \sum_{k=1}^N \left[\sqrt{1 + 16 h^2 \sin^2 \frac{p_k}{2}} - 1 \right] = \sum_{n=1}^{\infty} \gamma_{L, 2n}^{\text{CS}}(N) h^{2n}. \quad (2.5)$$

From the results of [41], we know the closed form of the *asymptotic* anomalous dimension at three orders for both twist 1 and 2. We recall here the expressions for the reader's benefit. With the usual definition of (nested) harmonic sums

$$S_a(N) = \sum_{n=1}^N \frac{(\text{sign } a)^n}{n^{|a|}}, \quad S_{a,b,\dots}(N) = \sum_{n=1}^N \frac{(\text{sign } a)^n}{n^{|a|}} S_{b,\dots}(n). \quad (2.6)$$

we have for twist 1

$$\gamma_{1,2}^{\text{CS}}(N) = 4 (S_1 - S_{-1}), \quad (2.7)$$

$$\gamma_{1,4}^{\text{CS}}(N) = -16(S_{-3} - S_3 + S_{-2,-1} - S_{-2,1} + S_{-1,-2} - S_{-1,2} - S_{1,-2} + S_{1,2} - S_{2,-1} + S_{2,1} + S_{-1,-1,-1} - S_{-1,-1,1} - S_{1,-1,-1} + S_{1,-1,1}), \quad (2.8)$$

and a much longer expression for $\gamma_{1,6}^{\text{CS}}(N)$ which can be found in [41]. All harmonic sums have in this case argument $S_a \equiv S_a(N)$. For twist-2, we find instead the compact expressions

$$\gamma_{2,2}^{\text{CS}}(N) = 4 S_1, \quad (2.9)$$

$$\gamma_{2,4}^{\text{CS}}(N) = 4 S_3 - 8 S_{1,2} - 4 S_{2,1}, \quad (2.10)$$

$$\gamma_{2,6}^{\text{CS}}(N) = 8 S_5 - 24 S_{1,4} - 32 S_{2,3} - 20 S_{3,2} - 16 S_{4,1} + 32 S_{1,1,3} + 24 S_{1,2,2} + 24 S_{1,3,1} + 20 S_{2,1,2} + 24 S_{2,2,1} + 8 S_{3,1,1} - 16 S_{1,1,2,1}. \quad (2.11)$$

and this time the argument is $S_a \equiv S_a(N/2)$.

These expressions do not include wrapping corrections. We shall compute the virtual scaling function from these asymptotic expressions assuming, as in $\mathcal{N} = 4$ SYM that it is independent on wrapping.

3. The fourth order twist-2 anomalous dimension

As remarked in [41] and in close analogy with twist-3 fields in $\mathcal{N} = 4$ SYM, the twist-2 anomalous dimensions involve nested harmonic sums with positive indices only. Such a restricted maximal transcendentality Ansatz reduces a lot the complexity of the determination of higher orders. In particular, we have determined the 8-th loop asymptotic anomalous dimension where dressing first appear. This will be a useful ingredient for the check of our results for the virtual scaling function. After a straightforward computation, we find

$$\gamma_{2,8}^{\text{CS}}(N) = G_{2,8}^{\text{CS}}(N) + \zeta_3 G_{2,8}^{\text{CS,dressing}}(N), \quad (3.1)$$

where

$$\begin{aligned}
G_{2,8}^{\text{CS}} = & 4(5 S_7 - 20 S_{1,6} - 39 S_{2,5} - 45 S_{3,4} - 35 S_{4,3} - 27 S_{5,2} - 15 S_{6,1} + 48 S_{1,1,5} + \\
& + 62 S_{1,2,4} + 58 S_{1,3,3} + 52 S_{1,4,2} + 38 S_{1,5,1} + 58 S_{2,1,4} + 64 S_{2,2,3} + \\
& + 68 S_{2,3,2} + 55 S_{2,4,1} + 44 S_{3,1,3} + 54 S_{3,2,2} + 52 S_{3,3,1} + 31 S_{4,1,2} + 33 S_{4,2,1} + \\
& + 26 S_{5,1,1} - 48 S_{1,1,1,4} - 52 S_{1,1,2,3} - 60 S_{1,1,3,2} - 52 S_{1,1,4,1} - 40 S_{1,2,1,3} + \\
& - 58 S_{1,2,2,2} - 58 S_{1,2,3,1} - 32 S_{1,3,1,2} - 42 S_{1,3,2,1} - 32 S_{1,4,1,1} - 32 S_{2,1,1,3} + \\
& - 54 S_{2,1,2,2} - 54 S_{2,1,3,1} - 38 S_{2,2,1,2} - 52 S_{2,2,2,1} - 42 S_{2,3,1,1} - 18 S_{3,1,1,2} + \\
& - 32 S_{3,1,2,1} - 34 S_{3,2,1,1} - 12 S_{4,1,1,1} + 32 S_{1,1,1,2,2} + 32 S_{1,1,1,3,1} + 16 S_{1,1,2,1,2} + \\
& + 32 S_{1,1,2,2,1} + 24 S_{1,1,3,1,1} + 4 S_{1,2,1,1,2} + 24 S_{1,2,1,2,1} + 28 S_{1,2,2,1,1} + \\
& + 20 S_{2,1,1,2,1} + 28 S_{2,1,2,1,1} + 12 S_{2,2,1,1,1} - 16 S_{1,1,1,2,1,1}), \tag{3.2}
\end{aligned}$$

$$G_{2,8}^{\text{CS,dressing}} = 32 (S_{2,2} + S_{3,1} - S_{1,2,1} - 2 S_{2,1,1}). \tag{3.3}$$

4. The virtual scaling function in ABJM from weak coupling

Up to vanishing terms for $N \rightarrow \infty$, the large spin expansion of our exact anomalous dimensions can be computed by the tricks reported in Appendix (A). Partial results can be found in [41]. We add here the 6 loop twist-1 and 8 loop twist-2 results. They read

$$\gamma_{L=1}^{\text{CS}}(N, h) = f^{\text{CS}}(h) (\log N + \gamma_{\text{E}} + \log 2) + B_{L=1}^{\text{CS}}(h) + \dots, \tag{4.1}$$

$$\gamma_{L=2}^{\text{CS}}(N, h) = f^{\text{CS}}(h) (\log N + \gamma_{\text{E}} - \log 2) + B_{L=2}^{\text{CS}}(h) + \dots, \tag{4.2}$$

where dots denote terms vanishing for large spin. The CS scaling function is half the $\mathcal{N} = 4$ value

$$f^{\text{CS}}(h) = \frac{1}{2} f^{\mathcal{N}=4}(h), \tag{4.3}$$

$$\begin{aligned}
f^{\mathcal{N}=4}(g) = & 8 g^2 - \frac{8}{3} \pi^2 g^4 + \frac{88}{45} \pi^4 g^6 - 16 \left(\frac{73}{630} \pi^6 + 4 \zeta_3^2 \right) g^8 + \\
& + 32 \left(\frac{887}{14175} \pi^8 + \frac{4}{3} \pi^2 \zeta_3^2 + 40 \zeta_3 \zeta_5 \right) g^{10} + \dots. \tag{4.4}
\end{aligned}$$

The virtual scaling function derived from harmonic sums turns out to be

$$B_{L=1}^{\text{CS}} = -12 \zeta_3 h^4 + \left(\frac{8 \pi^2 \zeta_3}{3} + 80 \zeta_5 \right) h^6 + \mathcal{O}(h^8), \tag{4.5}$$

$$B_{L=2}^{\text{CS}} = 4 \zeta_3 h^4 - 88 \zeta_5 h^6 + \left(-\frac{4}{15} \pi^4 \zeta_3 + 16 \pi^2 \zeta_5 + 1140 \zeta_7 \right) h^8 + \mathcal{O}(h^{10}) \tag{4.6}$$

These values must be compared with the known weak-coupling expansion valid for twist-

L fields in $\mathcal{N} = 4$ [45, 25, 46]

$$\begin{aligned}
B_L^{\mathcal{N}=4}(g) = & 8(2L - 7)\zeta_3 g^4 + \left(-\frac{8}{3}(L - 4)\pi^2 \zeta_3 - 8(21L - 62)\zeta_5\right) g^6 + \\
& + \left(\frac{8}{15}(3L - 13)\pi^4 \zeta_3 + \frac{8}{3}(11L - 32)\pi^2 \zeta_5 + 40(46L - 127)\zeta_7\right) g^8 + \\
& + \left(-64(2L - 7)\zeta_3^3 - \frac{128}{945}(11L - 49)\pi^6 \zeta_3 - \frac{8}{45}(103L - 310)\pi^4 \zeta_5 + \right. \\
& \left. - \frac{40}{3}(25L - 64)\pi^2 \zeta_7 - 392(55L - 146)\zeta_9\right) g^{10} + \dots .
\end{aligned} \tag{4.7}$$

5. The NLO BES equation in $\mathcal{N} = 4$

Let us recall how $B_L^{\mathcal{N}=4}$ is computed. In the by now standard notation of the BES paper [16], one has to solve the integral equation for the quantity $\sigma(t)$ closely related to the Fourier transform of the Bethe root density and obeying

$$\begin{aligned}
\sigma(t) = & \frac{t}{e^t - 1} \left[K(2gt, 0)(\log N + \gamma_E - (L - 2)\log 2) - \frac{L}{8g^2 t} (J_0(2gt) - 1) + \right. \\
& \frac{1}{2} \int_0^\infty dt' \left(\frac{2}{e^{t'} - 1} - \frac{L - 2}{e^{t'/2} + 1} \right) (K(2gt, 2gt') - K(2gt, 0)) + \\
& \left. - 4g^2 \int_0^\infty dt' K(2gt, 2gt') \sigma(t') \right].
\end{aligned} \tag{5.1}$$

The anomalous dimension is simply given by

$$\gamma_L^{\mathcal{N}=4}(g) = 16g^2 \sigma(0). \tag{5.2}$$

The kernel appearing in the above equation is

$$K(t, t') = K_0(t, t') + K_1(t, t') + K_d(t, t'), \tag{5.3}$$

with

$$K_0(t, t') = \frac{2}{tt'} \sum_{n \geq 1} (2n - 1) J_{2n-1}(t) J_{2n-1}(t'), \tag{5.4}$$

$$K_1(t, t') = \frac{2}{tt'} \sum_{n \geq 1} (2n) J_{2n}(t) J_{2n}(t'), \tag{5.5}$$

$$K_d(t, t') = 8g^2 \int_0^\infty dt'' K_1(t, 2gt'') \frac{t''}{e^{t''} - 1} K_0(2gt'', t'). \tag{5.6}$$

Now, the crucial point is that Eq. (5.1) is in close relation with the general properties of the transfer matrix which appear in the Baxter equation of $\mathcal{N} = 4$ as discussed in [45]. Since we want to derive a suitable modification of Eq. (5.1) valid in ABJM, we now analyze the Baxter equation for the ABJM twist operators.

6. Analysis of the ABJM $\mathfrak{sl}(2)$ Baxter equation

The following discussion will be done at the one-loop level will turn to be enough for our purposes. It is the adaptation of the methods described in [47] to the ABJM (twisted) case. The starting point is the Baxter $\widehat{Q}(u)$ -operator where u is the spectral parameter. It acts on the L -sites long $\mathfrak{sl}(2)$ spin chain and obeys the (twisted) Baxter equation [48]

$$\left(u + \frac{i}{2}\right)^L \widehat{Q}(u + i) - \left(u - \frac{i}{2}\right)^L \widehat{Q}(u - i) = \widehat{t}(u) \widehat{Q}(u). \quad (6.1)$$

The operator $\widehat{t}(u)$, also known as the auxiliary transfer matrix, is a polynomial of degree $L - 1$ in u with coefficients being local charges $\widehat{q}_n$ also acting on the spin chain states. The more precise form of $\widehat{t}(u)$ is

$$\widehat{t}(u) = i(2N + L)u^{L-1} + \sum_{n=2}^L \widehat{q}_n u^{L-n}. \quad (6.2)$$

Both the Baxter operator and the transfer matrix can be simultaneously diagonalized with the Hamiltonian. Replacing the operators with their eigenvalues, we obtain the same difference equation for the Baxter function $Q(u)$. The suitable boundary condition is simply the requirement that $Q(u)$ is a polynomial of degree N

$$Q(u) = \mathcal{N} \prod_{n=1}^N (u - u_n). \quad (6.3)$$

Its roots are readily identified with the Bethe roots since they obey the Bethe Ansatz equations. Notice that the operator $\widehat{Q}$ is required in order to compute the precise form of the energy eigenstates. Instead, for the eigenvalues, the polynomial $Q(u)$ is enough.

In practice, given a certain spin N , we can replace in the Baxter equation a polynomial Ansatz with undetermined charged q_n . Solving for the polynomial coefficients, we end with a set of algebraic equations for the charges. Each solution is associated with a specific $\mathfrak{sl}(2)$ module appearing in the subset of $[-1/2]^{\otimes L}$ states compatible with the twisted boundary conditions. The solutions δ_n of $t(\delta_n) = 0$ are called *holes*. They are dual solutions of the Bethe equations whose role will be clarified in a moment.

If we look for an even $Q(-u) = Q(u)$, then the transfer matrix (eigenvalue) has also a definite parity $t(-u) = (-1)^{L-1} t(u)$. Thus, the cases $L = 1, 2$ are particularly simple. For $L = 1$ we do not have holes nor charges. For $L = 2$ we have a single null hole $\delta = 0$ and, again, no charges. For $L > 2$, non-trivial charges start to appear. As discussed in [47, 45], the behaviour of the spin chain in the regime $N \gg L \gg 1$ is largely determined by the N dependence of the holes. In particular, it is important to determine whether the holes grow large as $N \rightarrow \infty$ or vanish. In $\mathcal{N} = 4$, it is known that two holes have a size which is $\mathcal{O}(N)$ while the other $L - 2$ holes are small in the sense that they vanish like $1/\log N$.

We claim that in the ABJM case, *all holes are $\mathcal{O}(1/\log N)$* . This statement can be efficiently checked by the methods of [47] which provide a good approximation to the holes

(accurate enough for our purposes) in the $N \gg L \gg 1$ regime. The main idea is that in the full region $u = \mathcal{O}((2N + L)^0)$ one can solve separately the two half-Baxter equations

$$\left(u + \frac{i}{2}\right)^L Q_+(u + i) = +t(u) Q_+(u), \quad (6.4)$$

$$\left(u - \frac{i}{2}\right)^L Q_-(u - i) = -t(u) Q_-(u). \quad (6.5)$$

where we write

$$t(u) = i(2N + L) \prod_{i=n}^{L-1} (u - \delta_n). \quad (6.6)$$

The solution is

$$Q_+(u) = [(2N + L)i]^{-iu} \frac{1}{\Gamma(1/2 - iu)^L} \prod_{n=1}^{L-1} \Gamma(i\delta_n - iu) \quad (6.7)$$

with Q_- being related to Q_+ by conjugation, regarding u as real. The approximate asymptotic Baxter function is then simply

$$Q^{\text{as}}(u) = Q_+(u) Q_-(-i/2) + Q_-(u) Q_+(i/2). \quad (6.8)$$

The request of zero residue at the holes leads to the quantization condition

$$\delta_n \log(2N + L) + L \text{Arg} \Gamma\left(\frac{1}{2} - i\delta_n\right) + \sum_{j=1}^{L-1} \text{Arg} \Gamma(1 + i\delta_n - i\delta_j) = \frac{\pi}{2} k_n, \quad (6.9)$$

where k_n are integers. Asymptotically this means

$$\delta_n = \frac{\pi}{2} \frac{k_n}{\log(N + L/2) + (2L + 1) \log 2 + \gamma_{\mathbb{E}}} = \mathcal{O}(1/\log N). \quad (6.10)$$

We can test the accuracy of Eq. (6.9) by comparing its solutions with the exact holes $\{\delta_n^{\text{exact}}\}$. The results are shown in the Tables reported in Tab. (1-2) for the non-trivial cases $L = 3, 4, 5$. The agreement is very good. We thus have very strong indications that all the $L - 1$ holes are vanishing for large spin. As an immediate (one-loop) consequence, we can evaluate the anomalous dimension which (from Eq. (6.8)) is

$$\gamma_{L,2}^{\text{CS}}(N) = 4 \log(2N + L) - 4\psi(1) + 2 \sum_{n=1}^{L-1} \left[\psi\left(\frac{1}{2} + i\delta_n\right) + \psi\left(\frac{1}{2} - i\delta_n\right) - 2\psi(1) \right], \quad (6.11)$$

where the second $-4\psi(1)$ term comes from the different numbers of factors in the last two terms of Eq. (6.7) and is absent in $\mathcal{N} = 4$. Taking the large N limit with $\delta_n \rightarrow 0$ we find

$$\gamma_{L,2}^{\text{CS}}(N) = 4 [\log N + \gamma_{\mathbb{E}} + (3 - 2L) \log 2] + \dots \quad (6.12)$$

In particular, this is in agreement with the explicit one-loop results at $L = 1, 2$.

7. The integral equation determining $B_L(g)$

From the results of the previous section, we can repeat the steps leading to Eq. (5.1). The main differences are listed below.

1. In $\mathcal{N} = 4$ SYM, the natural universal constants accompanying $\log N$ were the combination

$$\log N + \gamma_E - (L - 2) \log 2. \quad (7.1)$$

This is replaced in ABJM by the (one-loop) combination

$$\log N + \gamma_E - (2L - 3) \log 2. \quad (7.2)$$

2. The explicit factor $L - 2$ in Eq. (5.1) can be traced back to the number of small holes N_h which vanish for large N . It becomes $L - 1$ in ABJM. Similarly, there are various terms with factors 2 which are really $L - N_h$. This is 1 in ABJM.

In conclusion, we can write a modified Eq. (5.1) depending on a parameter ξ such that

$$\xi^{\mathcal{N}=4} = 1, \quad \xi^{\text{CS}} = \frac{1}{2}. \quad (7.3)$$

It reads

$$\begin{aligned} \sigma_\xi(t) = & \frac{t}{e^t - 1} \left[\xi K(2gt, 0) \left(\log \frac{N}{\xi} + \gamma_E - \left(\frac{L}{\xi} - 2 \right) \log 2 \right) - \frac{L}{8g^2 t} (J_0(2gt) - 1) + \right. \\ & \frac{1}{2} \int_0^\infty dt' \left(\frac{2\xi}{e^{t'} - 1} - \frac{L - 2\xi}{e^{t'/2} + 1} \right) (K(2gt, 2gt') - K(2gt, 0)) + \\ & \left. - 4g^2 \int_0^\infty dt' K(2gt, 2gt') \sigma_\xi(t') \right]. \end{aligned} \quad (7.4)$$

From this equation we immediately obtain the following exact scaling relation valid at NLO in the large spin expansion

$$\gamma_L^{\text{CS}}(N) = \frac{1}{2} \gamma_2^{\mathcal{N}=4}(2N). \quad (7.5)$$

As a corollary, we obtain Eq. (1.5) which is indeed verified by our explicit data at $L = 1, 2$.

8. Comments

In this paper, we have proved the scaling relation Eq. (1.4). This is not a totally surprising result. Indeed, it already appeared in the discussion of a rigid circular string stretched in both AdS_4 and along an S^1 of $\mathbb{CP}^3$ and carrying two spins [38]. It has also a counterpart in the case of the generalized scaling function of [45] as remarked in [33]. Here, we have given a proof valid in the case of twist operators belonging to a special $\mathfrak{sl}(2)$ subsector with many similarities to the $\mathcal{N} = 4$ SYM case. We have studied in details the necessary changes in the derivation of the integral equation for B_L^{CS} . This required the analysis of the asymptotic properties of the *hole* solutions to the twisted Baxter equation describing

this sector of ABJM. Eq. (1.4) is checked against the exact known anomalous dimensions of twist operators in ABJM presented in [41] as well as the new fourth order result for twist-2 illustrated here. Notice also that Eq. (7.5) is valid at all orders in the coupling. In particular it permits to obtain the strong coupling expansion of the virtual scaling function from the recent results of [25] with no effort.

As a final comment, let us remark that Eq. (1.4) answers a technical problem, *i.e.* the role of the phase deformation of ABJM at the level of the virtual scaling function. This is certainly interesting, but deserves a more sound physical motivation. We mentioned in the Introduction the important recent developments connecting the virtual scaling function of $\mathcal{N} = 4$ SYM to the properties of the on-shell scattering amplitudes. This connection has not been explored in ABJM. Compared to $\mathcal{N} = 4$ SYM, the quite different nature of the gauge theory side sets itself against the close integrability structure and, in our opinion, makes the problem an intriguing one.

Acknowledgments

We thank I. Klebanov, N. Gromov, D. Fioravanti, M. Rossi, and D. Bombardelli for comments.

A. Next-to-leading large spin expansion of harmonic sums

Let us remind a simple trick which reduces the NLO expansion of harmonic sums to the evaluation of multiple ζ values. The general large N expansion of a nested sum $S_a(N)$ has a singular part which is a polynomial in $\log N$, a constant term, and a remainder which vanishes at $N = \infty$. The degree of the singular polynomial equals the number of leading 1 indices in $\mathbf{a} = (a_1, \dots, a_\ell)$. This singular part can be extracted by using the shuffle algebra described in [49]. The main point is that if we start with ($a_1 \neq 1$, but it can also be absent)

$$S_{\underbrace{1, \dots, 1}_k, a_1, a_2, \dots, 1} \quad (\text{A.1})$$

we can add and subtract the product

$$\frac{1}{k+1} S_1 S_{\underbrace{1, \dots, 1}_k, a_1, a_2, \dots, 1}. \quad (\text{A.2})$$

Using the shuffle algebra in one of this terms, we cancel the initial sum in Eq. (A.1). Repeating iteratively this procedure we obtain the desired result. As an example let us consider $S_{1,1,1,2}$. Applying the above algorithm, we prove that

$$\begin{aligned} S_{1,1,1,2} = & \frac{1}{6} S_2 S_1^3 + \left(\frac{S_3}{2} - \frac{S_{2,1}}{2} \right) S_1^2 + \left(\frac{S_4}{2} - S_{3,1} + S_{2,1,1} \right) S_1 + \\ & + \frac{S_5}{6} + \frac{S_{2,3}}{3} - \frac{S_{3,2}}{6} - \frac{S_{4,1}}{2} + \frac{1}{2} S_{2,1,2} + S_{3,1,1} - S_{2,1,1,1}. \end{aligned} \quad (\text{A.3})$$

Taking the values at infinity we simply get

$$S_{1,1,1,2}(N) = \frac{1}{36}\pi^2 L^3 - \frac{1}{2}\zeta_3 L^2 + \frac{1}{40}\pi^4 L - \zeta_5 - \frac{1}{36}\pi^2 \zeta_3 + \dots, \quad (\text{A.4})$$

where $L = \log N + \gamma_E$ is the NLO expansion of $S_1(N)$.

References

- [1] N. Beisert and M. Staudacher, *Long-range PSU(2, 2|4) Bethe ansatz for gauge theory and strings*, Nucl. Phys. B **727**, 1 (2005) [arXiv:hep-th/0504190].
- [2] J. M. Maldacena, *The large N limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [3] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *A semi-classical limit of the gauge/string correspondence*, Nucl. Phys. B **636**, 99 (2002) [arXiv:hep-th/0204051].
- [4] S. Frolov and A. A. Tseytlin, *Semiclassical quantization of rotating superstring in AdS₅ × S⁵*, JHEP **0206**, 007 (2002) [arXiv:hep-th/0204226].
- [5] G. P. Korchemsky and G. Marchesini, *Structure function for large x and renormalization of Wilson loop*, Nucl. Phys. B **406**, 225 (1993) [arXiv:hep-ph/9210281].
- [6] A. V. Belitsky, A. S. Gorsky and G. P. Korchemsky, *Gauge / string duality for QCD conformal operators*, Nucl. Phys. B **667**, 3 (2003) [arXiv:hep-th/0304028].
- [7] A. V. Belitsky, A. S. Gorsky and G. P. Korchemsky, *Logarithmic scaling in gauge / string correspondence*, Nucl. Phys. B **748**, 24 (2006) [arXiv:hep-th/0601112].
- [8] L. F. Alday and J. M. Maldacena, *Comments on operators with large spin*, JHEP **0711**, 019 (2007) [arXiv:0708.0672 [hep-th]].
- [9] Yu. L. Dokshitzer, G. Marchesini and G. P. Salam, *Revisiting parton evolution and the large-x limit*, Phys. Lett. B **634**, 504 (2006) [arXiv:hep-ph/0511302].
- [10] Yu. L. Dokshitzer and G. Marchesini, *N = 4 SUSY Yang-Mills: Three loops made simple(r)*, Phys. Lett. B **646**, 189 (2007) [arXiv:hep-th/0612248].
- [11] B. Basso and G. P. Korchemsky, *Anomalous dimensions of high-spin operators beyond the leading order*, Nucl. Phys. B **775**, 1 (2007) [arXiv:hep-th/0612247].
- [12] M. Beccaria, V. Forini, T. Lukowski and S. Zieme, *Twist-three at five loops, Bethe Ansatz and wrapping*, arXiv:0901.4864 [hep-th].
- M. Beccaria and V. Forini, *Four loop reciprocity of twist two operators in N=4 SYM*, JHEP **0903**, 111 (2009) [arXiv:0901.1256 [hep-th]].
- M. Beccaria and V. Forini, *Reciprocity of gauge operators in N=4 SYM*, JHEP **0806**, 077 (2008) [arXiv:0803.3768 [hep-th]].
- M. Beccaria, *Three loop anomalous dimensions of twist-3 gauge operators in N=4 SYM*, JHEP **0709**, 023 (2007) [arXiv:0707.1574 [hep-th]].
- M. Beccaria, Yu. L. Dokshitzer and G. Marchesini, *Twist 3 of the sl(2) sector of N=4 SYM and reciprocity respecting evolution*, Phys. Lett. B **652**, 194 (2007) [arXiv:0705.2639 [hep-th]].
- M. Beccaria, *Universality of three gaugino anomalous dimensions in N = 4 SYM*, JHEP **0706**, 054 (2007) [arXiv:0705.0663 [hep-th]].
- M. Beccaria, *Anomalous dimensions at twist-3 in the sl(2) sector of N = 4 SYM*, JHEP **0706**, 044 (2007) [arXiv:0704.3570 [hep-th]].

- [13] M. Beccaria, V. Forini, A. Tirziu and A. A. Tseytlin, *Structure of large spin expansion of anomalous dimensions at strong coupling*, Nucl. Phys. B **812**, 144 (2009) [arXiv:0809.5234 [hep-th]].
- [14] Z. Bern, M. Czakon, L. J. Dixon, D. A. Kosower and V. A. Smirnov, *The Four-Loop Planar Amplitude and Cusp Anomalous Dimension in Maximally Supersymmetric Yang-Mills Theory*, Phys. Rev. D **75**, 085010 (2007) [arXiv:hep-th/0610248].
- [15] F. Cachazo, M. Spradlin and A. Volovich, *Four-Loop Cusp Anomalous Dimension From Obstructions*, Phys. Rev. D **75**, 105011 (2007) [arXiv:hep-th/0612309].
- [16] N. Beisert, B. Eden and M. Staudacher, *Transcendentality and crossing*, J. Stat. Mech. **0701**, P021 (2007) [arXiv:hep-th/0610251].
- [17] B. Basso, G. P. Korchemsky and J. Kotanski, *Cusp anomalous dimension in maximally supersymmetric Yang-Mills theory at strong coupling*, Phys. Rev. Lett. **100**, 091601 (2008) [arXiv:0708.3933 [hep-th]].
- [18] I. Kostov, D. Serban and D. Volin, *Functional BES equation*, JHEP **0808**, 101 (2008) [arXiv:0801.2542 [hep-th]].
- [19] L. J. Dixon, L. Magnea and G. Sterman, *Universal structure of subleading infrared poles in gauge theory amplitudes*, JHEP **0808**, 022 (2008) [arXiv:0805.3515 [hep-ph]].
- [20] A. H. Mueller, *On The Asymptotic Behavior Of The Sudakov Form-Factor*, Phys. Rev. D **20**, 2037 (1979).
J. C. Collins, *Sudakov form factors*, Adv. Ser. Direct. High Energy Phys. **5**, 573 (1989) [arXiv:hep-ph/0312336].
G. Sterman and M. E. Tejeda-Yeomans, *Multi-loop amplitudes and resummation*, Phys. Lett. B **552**, 48 (2003) [arXiv:hep-ph/0210130].
- [21] Z. Bern, L. J. Dixon and V. A. Smirnov, *Iteration of planar amplitudes in maximally supersymmetric Yang-Mills theory at three loops and beyond*, Phys. Rev. D **72**, 085001 (2005) [arXiv:hep-th/0505205].
- [22] F. Cachazo, M. Spradlin and A. Volovich, *Four-Loop Collinear Anomalous Dimension in $N = 4$ Yang-Mills Theory*, Phys. Rev. D **76**, 106004 (2007) [arXiv:0707.1903 [hep-th]].
- [23] G. Altarelli, *Partons In Quantum Chromodynamics*, Phys. Rept. **81**, 1 (1982).
- [24] L. F. Alday, *Universal structure of subleading infrared poles at strong coupling*, arXiv:0904.3983 [hep-th].
- [25] L. Freyhult and S. Zieme, *The virtual scaling function of AdS/CFT*, arXiv:0901.2749 [hep-th].
- [26] Z. Bajnok, R. A. Janik and T. Lukowski, *Four loop twist two, BFKL, wrapping and strings*, arXiv:0811.4448 [hep-th].
Z. Bajnok and R. A. Janik, *Four-loop perturbative Konishi from strings and finite size effects for multiparticle states*, Nucl. Phys. B **807**, 625 (2009) [arXiv:0807.0399 [hep-th]].
- [27] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, *$N=6$ superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, JHEP **0810**, 091 (2008) [arXiv:0806.1218 [hep-th]].
- [28] G. Arutyunov and S. Frolov, *Superstrings on $AdS_4 \times CP^3$ as a Coset Sigma-model*, JHEP **0809**, 129 (2008) [arXiv:0806.4940 [hep-th]].

- [29] B. . j. Stefanski, *Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$* Nucl. Phys. B **808**, 80 (2009) [arXiv:0806.4948 [hep-th]].
- [30] J. Gomis, D. Sorokin and L. Wulff, *The complete $AdS(4) \times CP(3)$ superspace for the type IIA superstring and D-branes*, JHEP **0903**, 015 (2009) [arXiv:0811.1566 [hep-th]].
- [31] J. A. Minahan and K. Zarembo, *The Bethe ansatz for superconformal Chern-Simons*, JHEP **0809**, 040 (2008) [arXiv:0806.3951 [hep-th]].
- [32] D. Bak and S. J. Rey, *Integrable Spin Chain in Superconformal Chern-Simons Theory*, JHEP **0810**, 053 (2008) [arXiv:0807.2063 [hep-th]].
- [33] N. Gromov and P. Vieira, *The all loop AdS_4/CFT_3 Bethe ansatz*, arXiv:0807.0777 [hep-th].
- [34] N. Gromov and P. Vieira, *The AdS_4/CFT_3 algebraic curve*, JHEP **0902**, 040 (2009) [arXiv:0807.0437 [hep-th]].
- [35] T. Nishioka and T. Takayanagi, *On Type IIA Penrose Limit and $N=6$ Chern-Simons Theories*, JHEP **0808**, 001 (2008) [arXiv:0806.3391 [hep-th]].
- [36] D. Gaiotto, S. Giombi and X. Yin, *Spin Chains in $N=6$ Superconformal Chern-Simons-Matter Theory*, arXiv:0806.4589 [hep-th].
- [37] G. Grignani, T. Harmark and M. Orselli, *The $SU(2) \times SU(2)$ sector in the string dual of $N=6$ superconformal Chern-Simons theory*, Nucl. Phys. B **810**, 115 (2009) [arXiv:0806.4959 [hep-th]].
- [38] T. McLoughlin, R. Roiban and A. A. Tseytlin, *Quantum spinning strings in $AdS_4 \times CP^3$: testing the Bethe Ansatz proposal*, JHEP **0811**, 069 (2008) [arXiv:0809.4038 [hep-th]].
- [39] C. Ahn and R. I. Nepomechie, *$N=6$ super Chern-Simons theory S-matrix and all-loop Bethe ansatz equations*, JHEP **0809**, 010 (2008) [arXiv:0807.1924 [hep-th]].
- [40] B. I. Zwiebel, *Two-loop Integrability of Planar $N=6$ Superconformal Chern-Simons Theory*, arXiv:0901.0411 [hep-th].
- [41] M. Beccaria and G. Macorini, *QCD properties of twist operators in the $\mathcal{N} = 6$ Chern-Simons theory*, arXiv:0904.2463 [hep-th].
- [42] T. McLoughlin and R. Roiban, *Spinning strings at one-loop in $AdS_4 \times P^3$* , JHEP **0812**, 101 (2008) [arXiv:0807.3965 [hep-th]].
L. F. Alday, G. Arutyunov and D. Bykov, *Semiclassical Quantization of Spinning Strings in $AdS_4 \times CP^3$* , JHEP **0811**, 089 (2008) [arXiv:0807.4400 [hep-th]].
C. Krishnan, *AdS_4/CFT_3 at One Loop*, JHEP **0809**, 092 (2008) [arXiv:0807.4561 [hep-th]].
- [43] S. A. Frolov, R. Roiban and A. A. Tseytlin, *Gauge - string duality for superconformal deformations of $N = 4$ super Yang-Mills theory*, JHEP **0507**, 045 (2005) [arXiv:hep-th/0503192].
- [44] N. Gromov and V. Mikhaylov, *Comment on the Scaling Function in $AdS_4 \times CP^3$* , JHEP **0904**, 083 (2009) [arXiv:0807.4897 [hep-th]].
- [45] L. Freyhult, A. Rej and M. Staudacher, *A Generalized Scaling Function for AdS/CFT* , J. Stat. Mech. **0807**, P07015 (2008) [arXiv:0712.2743 [hep-th]].
D. Fioravanti, G. Infusino and M. Rossi, *On the high spin expansion in the $sl(2)$ $\mathcal{N} = 4$ SYM theory*, arXiv:0901.3147 [hep-th].

- [46] D. Bombardelli, D. Fioravanti and M. Rossi, *Large spin corrections in $\mathcal{N} = 4$ SYM $sl(2)$: still a linear integral equation*, Nucl. Phys. B **810**, 460 (2009) [arXiv:0802.0027 [hep-th]].
D. Fioravanti, P. Grinza and M. Rossi, *Beyond cusp anomalous dimension from integrability*, arXiv:0901.3161 [hep-th].
- [47] G. P. Korchemsky, *Quasiclassical QCD pomeron*, Nucl. Phys. B **462**, 333 (1996) [arXiv:hep-th/9508025].
G. P. Korchemsky, *Integrable structures and duality in high-energy QCD*, Nucl. Phys. B **498**, 68 (1997) [arXiv:hep-th/9609123].
A. V. Belitsky, A. S. Gorsky and G. P. Korchemsky, *Logarithmic scaling in gauge / string correspondence*, Nucl. Phys. B **748**, 24 (2006) [arXiv:hep-th/0601112].
- [48] F. C. Alcaraz, M. N. Barber and M. T. Batchelor, *Conformal invariance, the XXZ chain and the operator content of two-dimensional critical systems*, Annals Phys. **182**, 280 (1988).
C. Korff, *A Q-operator for the twisted XXX model*, J. Phys. A: Math. Gen. **39**, 3203 (2006).
- [49] J. Blumlein, *Algebraic relations between harmonic sums and associated quantities*, Comput. Phys. Commun. **159**, 19 (2004) [arXiv:hep-ph/0311046].

N	$\delta_{L=3}^{\text{ex}}$	$\delta_{L=3}^{\text{qc}}$	$\delta_{L=4}^{\text{ex}}$	$\delta_{L=4}^{\text{qc}}$
4	0.22795472	0.22787539	0.40848291	0.40824222
8	0.21022561	0.2102162	0.37840139	0.37832836
12	0.20032517	0.20032661	0.3613073	0.36127663
16	0.1936300	0.19363366	0.3496743	0.34965917
20	0.18864721	0.18865113	0.3409907	0.34098254
24	0.18471752	0.1847212	0.33413129	0.33412668
28	0.18149529	0.18149861	0.32850149	0.32849884
32	0.17877816	0.17878113	0.32375148	0.32374999
36	0.17643811	0.17644076	0.31965923	0.31965845
40	0.17438935	0.17439172	0.31607559	0.31607526

Table 1: Comparison between the exact holes and the solutions to the quantization condition (qc) for $L = 3, 4$. In the first case, we have two opposite holes $\pm\delta$. In the second, we have three holes $0, \pm\delta$. We show the non trivial value δ in all cases. The integers k_n entering the quantization condition are ± 1 for $L=3$, and $\pm 2, 0$ for $L = 4$.

N	$\delta_{L=5}^{\text{ex}}$	$\delta_{L=5}^{\text{qc}}$	$(\delta_{L=5}^{\text{ex}})'$	$(\delta_{L=5}^{\text{qc}})'$
4	0.15014187	0.15013878	0.57240546	0.57206928
8	0.14425421	0.14424979	0.53079427	0.53066078
12	0.14046208	0.14045861	0.50680642	0.50673907
16	0.13769667	0.13769405	0.49041072	0.49037163
20	0.13553582	0.13553381	0.47815412	0.4781293
24	0.13377116	0.13376958	0.46846905	0.46845227
28	0.13228511	0.13228384	0.46052135	0.46050948
32	0.13100508	0.13100404	0.45381825	0.45380955
36	0.12988321	0.12988235	0.44804621	0.44803966
40	0.12888638	0.12888565	0.44299429	0.44298925

Table 2: Comparison between the exact holes and the solutions to the quantization condition (qc) for $L = 5$. The holes are $\pm\delta_{L=5}, \pm(\delta_{L=5})'$. We show the positive holes. The integers k_n entering the quantization condition are $\pm 3, \pm 1$.